A method for reductive cleavage of *N*-glycosylamide carbohydrate–peptide bond
L. M. LIKHOSHERSTOV, O. S. NOVIKOVA, V. E. PISKAREV, E. E. TRUSIKHINA, V. A. DEREVITSKAYA, AND N. K. KOCHETKOV (Moscow, U.S.S.R.) 155

The carbohydrate chains of influenza virus hemagglutinin
N. P. ARBATSKY, V. A. DEREVITSKAYA, A. O. ZHELTOVA, N. K. KOCHETKOV, L. M. LIKHOSHERSTOV, S. N. SENCHENKOVA, AND D. V. YURTOV (Moscow, U.S.S.R.) . . . 165

Studies on the chemical modification of soybean agglutinin
N. N. DESAI, A. K. ALLEN, AND A. NEUBERGER (London, Gt. Britain) 183

Structure of the capsular K3 antigen of *Escherichia coli* O4:K3:H4, a polysaccharide containing a 4-deoxy-2-hexulosonic acid
T. DENGLER, K. HIMMELSPACH, B. JANN, AND K. JANN (Freiburg, F.R.G.) 191

Effect of endoglycosidase F-peptidyl *N*-glycosidase F preparations on the surface components of the human erythrocyte
M. J. A. TANNER, D. J. ANSTEE, G. MALLINSON, K. RIDGWELL, P. G. MARTIN, N. D. AVENT, AND S. F. PARSONS, (Bristol, Gt. Britain) 203

Carbohydrate-binding specificities of five lectins that bind to *O*-glycosyl-linked carbohydrate chains. Quantitative analysis by frontal-affinity chromatography
S. SUEYOSHI, T. TSUJI, AND T. OSAWA (Tokyo, Japan) 213

Removal of ester-bound fatty acyl residues from *Salmonella* lipopolysaccharide by enzyme preparations from *Acanthamoeba castellanii*
W. DROZANSKI, A. E.-H. A. MOUSTAFA, O. LÜDERITZ, AND O. WESTPHAL (Freiburg, F.R.G.) 225

The comparison of monoclonal antibodies specific for two general epitopes located on *Citrobacter* O36 lipopolysaccharide
M. KULAKOWSKA, C. LUGOWSKI, E. ROMANOWSKA (Wrocław, Poland) AND D. R. BUNDLE (Ottawa, Ontario, Canada) 233

Immunochemical studies on the *N*-acetyllactosamine β-(1→6)linked trisaccharide specificity of *Ricinus communis* agglutinin
A. M. WU, S. SUGII (College Station, TX, U.S.A.), F. G. GRUEZO AND E. A. KABAT (New York, NY, U.S.A.) 243

Substrate recognition by UDP-*N*-acetyl-α-D-galactosamine: polypeptide *N*-acetyl-α-D-galactosaminyltransferase. Effects of chain length and disulphide bonding of synthetic peptide substrates
R. C. HUGHES, A. F. BRADBURY, AND D. G. SMYTH (London, Gt. Britain) 259

Blood group Tn-active macromolecules from human carcinomas and erythrocytes. Characterization of and specific reactivity with mono- and poly-clonal anti-Tn antibodies induced by various immunogens
G. F. SPRINGER, E. V. CHANDRASEKARAN, P. R. DESAI, AND H. TEGTMEYER (Evanston, IL, U.S.A.) 271

The binding of the Lewis-a human blood group determinant by two hybridoma monoclonal anti-Le^a antibodies
R. U. LEMIEUX, O. HINDSGAUL, P. BIRD, S. NARASIMHAN (Edmonton, Alberta, Canada), AND W. W. YOUNG, JR. (Charlottesville, VA, U.S.A.) 293

(Continued overleaf)

(contents continued)

The effect of glycosidases on the survival of rat erythrocytes in circulation
K. A. GUTOWSKI, D. A. LINSEMAN, AND D. AMINOFF (Ann Arbor, MI, U.S.A.) . . 307

Notes

Oxidation products of ovarian cyst mucins. Characterisation by density-gradient analysis
J. M. CREETH (Bristol, Gt. Britain) 314

Blood-group active proteoglycan containing 3-*O*-methylrhamnose (acofriose) from young plants of *Osmunda japonica*
T. AKIYAMA, K. TANAKA, S. YAMAMOTO, AND S. ISEKI (Tokyo, Japan) 320

Structural analysis of the preponderant high-mannose oligosaccharide of human Tamm-Horsfall glycoprotein
F. DALL'OLIO, F. J. J. DE KANTER, D. H. VAN DEN EIJNDEN, AND F. SERAFINI-CESSI (Amsterdam, The Netherlands) 327

Analysis of a mouse monoclonal antibody detecting the H Type 1 blood group determinant
S. WELT, B. W. T. YIN, K. FURUKAWA, K.-A. KARLSSON, L. J. OLD, AND K. O. LLOYD (Göteborg, Sweden) 333

Announcement C1

Author index C3

Subject index C5